
Enhanced paracrine FGF10 expression promotes formation of multifocal prostate adenocarcinoma and an increase in epithelial androgen receptor.

Journal: Cancer Cell

Publication Year: 2007

Authors: Sanaz Memarzadeh, Li Xin, David J Mulholland, Alka Mansukhani, Hong Wu, Michael A Teitell, Owen N Witte

PubMed link: 18068633

Funding Grants: Role of Mitochondria in Self-Renewal Versus Differentiation of Human Embryonic Stem Cells, CIRM Type I Comprehensive Training Program

Public Summary:

Scientific Abstract:

Enhanced mesenchymal expression of FGF10 led to the formation of multifocal PIN or prostate cancer. Inhibition of epithelial FGFR1 signaling using DN FGFR1 led to reversal of the cancer phenotype. A subset of the FGF10-induced carcinoma was serially transplantable. Paracrine FGF10 led to an increase in epithelial androgen receptor and synergized with cell-autonomous activated AKT. Our observations indicate that stromal FGF10 expression may facilitate the multifocal histology observed in prostate adenocarcinoma and suggest the FGF10/FGFR1 axis as a potential therapeutic target in treating hormone-sensitive or refractory prostate cancer. We also show that transient exposure to a paracrine growth factor may be sufficient for the initiation of oncogenic transformation.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/enhanced-paracrine-fgf10-expression-promotes-formation-multifocal-prostate>